

2024 Annual Report City of Fort Collins Mosquito Control Program



Vector Disease Control International 318 N. Garfield Ave Loveland, CO 80537 Phone 970-278-9977

Website: www.vdci.net/Colorado

City of Fort Collins Mosquito Management Operations

Annual Report For 2024

Table of Contents

| | Page |
|---|---------|
| PROGRAM OBJECTIVES | 3 |
| VDCI'S COMMITMENT | 3 |
| 2024 SEASON PERSPECTIVE CLIMATE COMPARISON DATA | 4 |
| WEST NILE VIRUS SEASON WV ACTIVITY BY STATE (CDC 2024 HUMAN CASE MAP) NEUROINVASIVE INCIDENCE BY STATE (CDC 2024 HUMAN CASE MAP) COLORADO WNUV DATA (CDPHE 2024 HUMAN CASE REPORTS) | 5 |
| LARVAL MOSQUITO CONTROL LARVAL SITE INSPECTIONS & TREATMENTS IN FORT COLLINS LARVAL ACREAGE TREATMENTS IN FORT COLLINS LARVICIDE PRODUCT APPLICATION BY TYPE | 7 |
| VDCI SURVEILLANCE LABORATORY CDC SURVEILLANCE LIGHT TRAP DATA ADULT MOSQUITO LIGHT TRAP GENUS SUMMARY 2024 CSU WEST NILE VIRUS MOSQUITO SAMPLING RESULTS – LARIMER COUNTY | 8 |
| ADULT MOSQUITO CONTROL SEASON DETAILS ADULT MOSQUITO CONTROL APPLICATION REPORT FOR FORT COLLINS 2024 | 12 1 |
| PUBLIC RELATIONS AND EDUCATION MOSQUITO LINE CALLS FOR FORT COLLINS | 13 |
| APPENDIX 1. FORT COLLINS INDIVIDUAL LIGHT TRAP SUMMARIES 2024 2. LARIMER COUNTY ADULT SAMPLE POOL POSITIVE RESULTS AND LOCATION | 15 |

Program Objectives

Vector Disease Control International, LLC (VDCI) completed its 21st year of cost-effective Integrated Mosquito Management (IMM) for Fort Collins in 2024. The primary objective of Fort Collins' IMM Program is to monitor and reduce mosquito populations through the use of specific, environmentally sound, control techniques in order to protect its residents from the threat of mosquito-borne diseases. VDCI prioritizes the detection and elimination of larval mosquitoes in aquatic habitats, in conjunction with the monitoring of adult mosquito populations through routine surveillance, in order to assess West Nile virus vector species abundance in the area.

Open communication is maintained by VDCI between Residents, Property Management Companies, the Weld and Larimer County Departments of Health & Environment and surrounding municipalities to ensure that the highest level of mosquito control and epizootic response is achieved. This diligent and cooperative communication is important to the Fort Collins mosquito management program and provides significant benefit to public health throughout the entire area.


VDCI's Commitment

Vector Disease Control International is a company built on the foundations of public health, ethics, professionalism, and technical expertise. VDCI is committed to providing our customers with scientifically based, environmentally sensitive and technologically advanced Integrated Mosquito Management (IMM) programs of the highest quality. All of our employees are committed to excellence in vector control and public health and strive to improve the quality of human life in communities through public education and the control of mosquitoes and the diseases they can transmit. VDCI currently has programs across the state of Colorado, providing services for towns, cities, counties, homeowners associations, and encephalitis surveillance monitoring programs for county health departments.


Vector Disease Control International, as the contractor for Fort Collins, will continue to use proven scientific Integrated Mosquito Management techniques to survey and control local mosquito populations using biorational larval controls and limited low-toxicity insecticide applications. All of the methods and materials used have been reviewed and registered by the US Environmental Protection Agency, the Centers for Disease Control, the Colorado Department of Agriculture and the American Mosquito Control Association.

2024 Season Perspective

At VDCI, we anticipate each Colorado summer to bring a unique combination of temperature, precipitation, irrigation, and human factors that create new challenges for mosquito control and mosquito-borne disease management. In 2024, Northern Colorado experienced a hot and dry year. Larimer and Weld counties, in particular, saw a significant moisture deficit, with some areas receiving nearly 43% less precipitation than usual. This reduced precipitation led to less standing water and lower mosquito numbers throughout the season. Overall, 2024 can be considered a 'light' mosquito season.



During the summer of 2024, Northern Colorado experienced hotter-than-normal temperatures. Both Larimer and Weld counties saw a 4-6°F increase from the average. This extra heat and evaporation further dried the region, leading to a decrease in mosquito proliferation due to the reduced standing water. In contrast to the record precipitation levels of 2023, this reduction in mosquito activity was a welcome change in 2024. Analysis of trap data from various locations in Larimer and Weld counties reveals an average of 82% fewer mosquitoes trapped in 2024 compared to 2023.




West Nile Virus Season

Since the introduction of West Nile virus to the United States in 1999, the virus has made a complete westward expansion to the West Coast. Starting in the Northeastern parts of the United States, the virus steadily spread through the South, the Midwest, the Rocky Mountain region, and to the Western States. This extensive distribution is due to the ability of WNv to establish and persist in the wide variety of ecosystems present across the country. WNv has been detected in 65 different mosquito species in the U.S., though it appears that only a few Culex species drive epizootic and epidemic transmission (WNv Guidelines, CDC 2013). Although West Nile virus has been endemic to the United States since 1999, researchers continue to seek an understanding for some of the factors which contribute to region specific spikes in vector abundance and human risk. We still do not understand why some humans develop West Nile fever while other infections develop into more serious West Nile encephalitis or West Nile meningitis cases. Additionally, physicians and researchers continue to seek answers to the variable recovery times and occurrence of deaths that result with some infections. WNv has expanded to the point that it can now be found in all 48 contiguous states and since its introduction has produced two additional, large nationwide epidemics in 2003 and 2012 (WNv Guidelines, CDC 2013).

As of September 17th, 2024, the CDC has reported West Nile virus infections in people in 43 states and the District of Columbia. Overall, 659 cases of West Nile virus disease in people have been reported to CDC. Of these, 452 (68.6%) were classified as neuroinvasive disease (such as meningitis or encephalitis) and 207 (31.4%) were classified as non-neuroinvasive disease.


West Nile virus human disease cases reported by state of residence, 2024



2024 Annual Report of Mosquito Management Operations Vector Disease Control International

Colorado 2024

As of September 11th, 2024 The Colorado Department of Health and Environment reports a total of 48 human cases of West Nile virus infection from the state of Colorado. Fifteen counties across the state of Colorado have reported human West Nile virus infection. In Northern Colorado, 3 human cases are reported from Larimer County, and 11 from Weld County. While we have passed the historical peak of WNv risk these numbers are expected to rise as there is often a delay in onset of symptoms, diagnosis and reporting. Please note that any additional cases reported by CDPHE will also be reported to the Centers for Disease Control.



People affected: 48

Affected people who died: 2

Affected people with neurologic symptoms: 24

Affected people requiring hospitalization: 16

2024 cases
Previous 5 yr average

Larval Mosquito Control

Larval mosquito control is the foundation of the Fort Collins Mosquito Control program and can be an extremely effective way to manage mosquitoes, thereby reducing the number of potential disease vectors and annoyances associated with biting adults. Years of research and practical experience have shown that the most effective way to control mosquito populations is through an aggressive Integrated Mosquito Management (IMM) approach. This approach aims at using a variety of concepts, tools, and products to reduce a pest population to a tolerable level.

Pre-season larval control work involved ground truthing GIS maps and remapping areas where new development or flooding had altered the landscape. VDCI began larval site inspections in many areas mid-April. Hiring of seasonal field technicians began in March and continued into May. VDCI's Annual Field Technician Classroom Training Day took place on May 28th. Field training by VDCI management and veteran employees lasted through May and full-time field activities were in full force by early June.

In 2024, Vector Disease Control field technicians performed 5,554 larval site inspections, of which 3,959 (71.4%) were wet upon inspection, 1,952 (49.3% of wet sites) were producing mosquito larvae. To prevent these larvae from emerging as adult mosquitoes, VDCI applied 5,992.5 lbs. of VectoBac (Bti) and 8.3 gal of BVA 2 mineral oil to 2,196.4 acres of land.

During 2024, VDCI technicians performed site inspections at 12 residences as part of our backyard inspection program. VDCI also performed treatments at approximately 650 stormdrains. Drains that were wet or had the potential to hold water were treated with long term larvicide products including Altosid pellets, briquettes, and VectoLex WSP (Bsph).



Larval mosquito control can be achieved in several ways including biological, biochemical, chemical, and mechanical means. No single larvicide product will work effectively in every habitat where mosquito larvae are found, so a variety of products and methods should be employed. Additionally, although there are a variety of methods for reducing larval populations, some may have negative consequences that outweigh their benefits. Mechanical or physical habitat modification is a technique which VDCI uses on relatively small scale projects, as the area to be modified must be carefully reviewed.


VDCI's favored method of larval mosquito control is through the use of bacterial bio-rational products. The main product used by VDCI is a variety of bacteria (*Bacillus thuringiensis var. israeliensis*). *Bti*, as it is known, has become the cornerstone of mosquito control programs throughout the world. The benefits include its efficacy and lack of environmental impacts. When used in accordance with its label, successful control of mosquito larvae can be achieved without impact to non-target species such as other aquatic invertebrates, birds, mammals, fish, amphibians, reptiles, or humans. A broad label allows for the use of the product in the

majority of the habitats throughout the service area. Another bacterial product closely related to *Bti* is *Bacillus sphaericus* (*Bs*). *BS* provides similar benefits to *Bti* while also providing residual control of certain species of mosquitoes. It is used specifically in difficult to treat areas where *Culex* are the predominant species due to its limitations and high cost.

Other larval control products include the insect growth regulator methoprene (Altosid), and light mineral oils (BVA 2 larvicide oil). Methoprene is a synthetic version of a juvenile growth hormone in larval mosquitoes. The hormone prevents the normal development of larval mosquitoes into pupae and adults, eventually causing death. VDCI limits the use of chemical larvicides to areas with little biodiversity, such as road side ditches, or areas that chronically produce high mosquito populations. They are only used after a thorough assessment has been made of any habitat where their use is being considered. Mineral oil is the only product effective in controlling mosquito pupae and therefore is an essential tool when pupae are present.

VDCI Surveillance Laboratory

Information about mosquito abundance and species diversity is essential to integrated program. Vector Disease Control International utilizes two kinds of traps to monitor mosquito populations. The most commonly used is the CDC light trap which uses carbon-dioxide from



dry ice as bait to attract female mosquitoes seeking a blood meal from a breathing animal. Once attracted by the CO_2 , the mosquitoes are lured by a small light to a fan that pulls them into a net for collection. The second type of trap VDCI uses is called a gravid trap. Gravid traps use a tub of highlyorganic water as bait to attract female mosquitoes that are looking for a place to lay their eggs. A fan placed close to the water surface forces mosquitoes that come to the water into a collection net. Once back in the laboratory, the contents of the trap nets are counted and speciated by trained technicians.

In 2024, Vector Disease Control International monitored a statewide network of hundreds

of weekly trap sites, collecting adult mosquitoes that were counted and identified to species by the VDCI Surveillance Laboratories. While individual traps provide only limited information, trap data is interpreted in the context of historical records for the same trap site, going back in time more than a decade. Individual traps are also compared to other traps from around the region that were set on the same night and therefore exposed to similar weather conditions. Technicians working in the Surveillance Laboratories at Vector Disease Control International are trained to provide accurate species-level identification of both larval and adult mosquitoes.

Additionally, the VDCI Surveillance Laboratory conducts an intensive larval identification program with larval mosquito samples collected by I&L technicians prior to larviciding being

identified to species. This information is now invaluable in targeting mosquito control efforts as we gain a greater understanding of the habitat types preferred by Colorado mosquito species and the seasonality of these habitats as sites for mosquito development.


Specimens and data collected from these traps and larval identification are used in:


- Determining the effect of larval control efforts. Each mosquito species prefers specific kinds of habitats for larval development. If a trap includes large numbers, it could indicate the presence of an unknown larval habitat and, based on the species identification and known habitat preference for that species, direct field technicians as to possible sources of the mosquitoes collected.
- <u>Determining larval and adult mosquito species.</u> This helps to illustrate the threat of mosquito-borne disease amplification and transmission because different mosquito species can vector different diseases to people and animals.
- Determining where adult control efforts were necessary. While mosquito eradication is impossible, significant population reduction is achievable. In places where larval control is insufficient, such as neighborhoods where adult mosquitoes have migrated in from outside of the control area, it may be necessary to use adulticide methods, such as ULV truck fogging or barrier sprays of harborage areas. Trap counts that exceed an acceptable threshold for an area may trigger adult control measures.
- Surveillance for Mosquito-borne Disease. Historically, VDCI efforts were targeted primarily at controlling mosquito nuisance problems with limited disease surveillance. However, since the arrival of the West Nile virus in Colorado in August of 2002, the paradigm has shifted toward disease prevention and control. Accurate species identification of the mosquitoes in the traps is important when monitoring species population trends. It also is necessary for evaluating whether a population spike represents an actual increase in disease transmission potential or only an increased nuisance level.

SURVEILLANCE LIGHT TRAP DATA

In 2024, there were 783 CDC light surveillance traps set within Fort Collins, which collected a total of 40,597 mosquitoes. This year there was an average of 51.8 mosquitoes caught per trap per night and an average 17.5 *Culex* mosquitoes per trap per night. The composition of mosquitoes collected was 33.7% (13,682) *Culex spp.*, 61.3% (24,882) *Aedes/Ochlerotatus spp.*, 1.1% (430) *Culiseta spp.*, and 1.3% (543) *Anopheles spp.*


2024 Fort Collins Trap Composite Data





Adult Mosquito Surveillance Summary

Start Date: 05/01/2024 End Date: 09/17/2024



CSU WEST NILE VIRUS MOSQUITO SAMPLE TESTING RESULTS - LARIMER COUNTY

Many local health departments have moved towards mosquito-based surveillance indicators to assess the weekly risk of West Nile transmission and guide response decisions for adult mosquito control applications. The vector index and infection rate is derived by testing the mosquitoes VDCI collects for the presence of West Nile virus. This value is closely monitored by the CDPHE and local health departments to evaluate the risk posed by the vector mosquito population.

As defined in the CDC guidelines for West Nile virus surveillance, prevention and control, the vector index (VI) is an estimate of the number of West Nile virus infected mosquitoes in an area. This number can serve as a human health risk value. An operational value of 0.5, which was derived from the comparison of historical data for human infections, as well as relative abundance and infection in mosquitoes, serves as an indicator of high risk for West Nile virus transmission to humans in the corresponding area. As the value of the vector index increases there is a corresponding risk of human disease and this value can be used to offset epidemics.

As stated on the CDPHE website, the seasonal variation of West Nile virus risk can change throughout a summer and it is best to assume you have some risk for WNV if you have mosquitoes.

As of Sept 16th, 2024 Colorado State University's Department of Microbiology, Immunology and Pathology, has tested a total of 1,414 mosquito pools submitted by VDCI that were collected from Larimer County. A total of 52 mosquito pool samples have tested positive for WNv with 2 of those being collected from Berthoud, 6 from Loveland and 44 from Fort Collins. Testing of these mosquitoes for West Nile virus is paid for by the Town of Berthoud, the City of Fort Collins, and the City of Loveland. It's important to note that the large number of WNv positive sample pools reported from Fort Collins is highly correlated with the fact they test all mosquitoes with the potential for transmitting disease versus just a subset of the population. Also important to note that in 2024, additional West Nile virus pool testing was conducted in Berthoud by Larimer County Public Health in conjunction with the CDC, the above numbers reflect only those submitted by VDCI.

ADULT MOSQUITO CONTROL

The goal of Vector Disease Control International is to provide our customers with the best options for safe, effective, modern mosquito management. The primary emphasis of the Fort Collins Mosquito Management Program is to control mosquitoes in the larval stage, using safe biological control products. VDCI uses EPA and CDC approved adulticides to reduce mosquito populations. There were no adult mosquito control operations contracted by the City of Fort Collins during 2024.

VDCI has been privately contracted to perform adult mosquito control for Countryside HOA, Greenstone HOA, Lindenwood HOA, and Willow Springs HOA, during the 2024 season. A total of 34.13 road miles within these neighborhoods were sprayed via ground based vehicles with Ultra Low Volume (ULV) AquaPermX 3030 mosquito adulticides to reduce adult mosquito populations on the dates listed below.

| Date | Zone | Spray Miles | Ounces Sprayed |
|-----------|----------------------------------|-------------|-------------------|
| 6/5/2024 | LHOA - Countryside Spray Zone | 1.47 | 57.07 |
| 6/5/2024 | LHOA - Willow Springs Spray Zone | 3.05 | 109.11 |
| 6/5/2024 | LHOA - Greenstone HOA Spray Zone | 2.63 | 92.54 |
| 6/19/2024 | LHOA - Lindenwood HOA Spray Zone | 2.21 | 79.85 |
| 6/26/2024 | LHOA - Willow Springs Spray Zone | 2.76 | 98.73 |
| 6/26/2024 | LHOA - Lindenwood HOA Spray Zone | 2.26 | 80.73 |
| 7/10/2024 | LHOA - Countryside Spray Zone | 1.66 | 62.36 |
| 7/17/2024 | LHOA - Willow Springs Spray Zone | 3.74 | 132.19 |
| 7/3/2024 | LHOA - Lindenwood HOA Spray Zone | 3.54 | 126.58 |
| 7/10/2024 | LHOA - Lindenwood HOA Spray Zone | 2.18 | 77.44 |
| 8/7/2024 | LHOA - Willow Springs Spray Zone | 3.79 | 135.93 |
| 8/7/2024 | LHOA - Countryside Spray Zone | 1.49 | 52.53 |
| 8/28/2024 | LHOA - Willow Springs Spray Zone | 3.35 | 127.41 |

VDCI uses state of the art technology, calibrated application timing, and least-toxic products to minimize non-target impacts. All adult mosquito control is accomplished using Ultra Low Volume (ULV) fogging equipment and performed after dusk when the majority of mosquito species are most active. This type of equipment produces droplets averaging 12 microns in diameter and allows for a minimal amount of product to be put into the environment. These treatments take place in the evening when mosquitoes are flying in greater numbers and nontarget insect activity (for example, day-flying pollinators like bees) is greatly reduced. Using this application technique, the overall goal of minimal environmental impact and effective adult control is achieved in the targeted area.

Public Relations and Education

VDCI is dedicated to providing strong Public Outreach and Education Programs to residents in all of our communities. Citizen complaints, inquiry, information and satisfaction surveys can aid in evaluating the effectiveness of a program. VDCI constantly looks for ways to better serve the communities we work with and encourages both the citizen and local media involvement in order to increase the effectiveness of our programs. We have clearly demonstrated that commitment and belief by proactively serving Fort Collins (and all of our contracted communities) with numerous innovative programs, activities and services.

Customer service is always a high priority for VDCI. We take pride in training each and every technician so that they have the knowledge to provide residents with the correct answers to their questions. Each field technician spends part of their day responding to resident concerns in their work area. This in-field customer service personalizes the mosquito control program, provides VDCI with local information on mosquito activity and presents a valuable opportunity to educate our residents about mosquito biology and control.

MosquitoLine™

VDCI maintains a toll-free telephone line (877-276-4306) and local line to the Northern Colorado Office (970-278-9977) at no cost to the customer. This service can be utilized to accept calls from the public concerning:

- Information about mosquito biology and source reduction of mosquito habitats
- information on program components, operations and monitoring
- Seasonal West Nile virus activity
- Personal protection options for mosquito annoyances and West Nile virus risk
- Reports about mosquitoes and possible larval mosquito habitats
- Requests to perform larvicide applications and/or opt-out of any adulticide spraying via a shut-off list
- Request notification when adulticide spraying is planned in their neighborhood

Request health and safety information about mosquito control operations and pesticide products used

VDCI has provided Mosquito Hotlines to the residents in communities which we are contracted to also reduce workload by municipal personnel. This enables direct communication and response by mosquito control employees to resident's concerns about West Nile virus and larval site activity and treatment. VDCI maintains a log of calls received and will summarize call activity in monthly and annual reports.

VDCI received 7 phone calls from Fort Collins residents in 2024. VDCI received 5 calls from residents reporting larval mosquito sites in their area. VDCI inspected all of the reported sites and added new sites if found producing mosquito larvae and were not already part of our regular inspection program. VDCI received 1 mosquito annoyance call from residents of the City of Fort Collins, who were requesting information on how the decisions on adulticiding are made. VDCI provided the relevant information and directed them to the city's website for further information. VDCI received 1 phone call from residents regarding any upcoming adult mosquito control applications. All relevant questions were answered, and the residents were guide to www.fcgov.com/westnile for further information.

Appendix 1: Individual Light Trap Summaries FC-001

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 Magic Carpet

GPS: 40.49087011315602, -105.08256990462542

 Total number of trap/nights set:
 15.0

 Total number of mosquitoes collected:
 222.0

 Average mosquitoes per trap/night:
 14.8

 Average Culex per trap/night:
 5.4

Species collected and abundance:

| Aedes dorsalis | 23.0 | 10.4% |
|-------------------|-------|-------|
| Aedes increpitus | 2.0 | 0.9% |
| Aedes vexans | 115.0 | 51.8% |
| Culex pipiens | 14.0 | 6.3% |
| Culex tarsalis | 67.0 | 30.2% |
| Culiseta inornata | 1.0 | 0.5% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 140.0 | 63.1% |
| Anopheles | 0.0 | 0.0% |
| Culex | 81.0 | 36.5% |
| Culiseta | 1.0 | 0.5% |
| Other | 0.0 | 0.0% |

FC-004

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 Big Horn

GPS: 40.53498867239214, -105.03751885145903


 Total number of trap/nights set:
 15.0

 Total number of mosquitoes collected:
 591.0

 Average mosquitoes per trap/night:
 39.4

 Average Culex per trap/night:
 28.3

| Aedes dorsalis | 57.0 | 9.6% |
|---------------------------|-------|-------|
| Aedes increpitus | 1.0 | 0.2% |
| Aedes melanimon | 2.0 | 0.3% |
| Aedes vexans | 98.0 | 16.6% |
| Coquillettidia perturbans | 2.0 | 0.3% |
| Culex pipiens | 156.0 | 26.4% |
| Culex tarsalis | 269.0 | 45.5% |
| Culiseta inornata | 6.0 | 1.0% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 158.0 | 26.7% |
| Anopheles | 0.0 | 0.0% |
| Culex | 425.0 | 71.9% |
| Culiseta | 6.0 | 1.0% |
| Other | 2.0 | 0.3% |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 N. Linden

GPS: 40.592327859908366, -105.06629493087529

 Total number of trap/nights set:
 15.0


 Total number of mosquitoes collected:
 134.0

 Average mosquitoes per trap/night:
 8.9

 Average Culex per trap/night:
 3.4

Species collected and abundance:

| Aedes dorsalis | 13.0 | 9.7% |
|---------------------------|------|-------|
| Aedes nigromaculis | 1.0 | 0.7% |
| Aedes vexans | 63.0 | 47.0% |
| Coquillettidia perturbans | 1.0 | 0.7% |
| Culex pipiens | 16.0 | 11.9% |
| Culex tarsalis | 35.0 | 26.1% |
| Culiseta inornata | 5.0 | 3.7% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 77.0 | 57.5% |
| Anopheles | 0.0 | 0.0% |
| Culex | 51.0 | 38.1% |
| Culiseta | 5.0 | 3.7% |
| Other | 1.0 | 0.7% |

FC-011

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

ocation: Golden Current

GPS: 40.56947993409379, -105.13603001832962


 Total number of trap/nights set:
 15.0

 Total number of mosquitoes collected:
 812.0

 Average mosquitoes per trap/night:
 54.1

 Average Culex per trap/night:
 6.3

| Aedes cinereus | 3.0 | 0.4% |
|---------------------|-------|-------|
| Aedes dorsalis | 2.0 | 0.2% |
| Aedes increpitus | 112.0 | 13.8% |
| Aedes melanimon | 3.0 | 0.4% |
| Aedes trivittatus | 1.0 | 0.1% |
| Aedes vexans | 571.0 | 70.3% |
| Anopheles freeborni | 8.0 | 1.0% |
| Culex pipiens | 56.0 | 6.9% |
| Culex tarsalis | 39.0 | 4.8% |
| Culiseta inornata | 17.0 | 2.1% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 692.0 | 85.2% |
| Anopheles | 8.0 | 1.0% |
| Culex | 95.0 | 11.7% |
| Culiseta | 17.0 | 2.1% |
| Other | 0.0 | 0.0% |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 FC Visitor Center

GPS: 40.56509442567909, -105.00627581030129

 Total number of trap/nights set:
 15.0


 Total number of mosquitoes collected:
 649.0

 Average mosquitoes per trap/night:
 43.3

 Average Culex per trap/night:
 26.4

Species collected and abundance:

| Aedes dorsalis | 92.0 | 14.2% |
|---------------------------|-------|-------|
| Aedes increpitus | 22.0 | 3.4% |
| Aedes melanimon | 11.0 | 1.7% |
| Aedes nigromaculis | 1.0 | 0.2% |
| Aedes trivittatus | 2.0 | 0.3% |
| Aedes vexans | 103.0 | 15.9% |
| Anopheles freeborni | 5.0 | 0.8% |
| Coquillettidia perturbans | 6.0 | 0.9% |
| Culex pipiens | 99.0 | 15.3% |
| Culex tarsalis | 297.0 | 45.8% |
| Culiseta inornata | 11.0 | 1.7% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 231.0 | 35.6% |
| Anopheles | 5.0 | 0.8% |
| Culex | 396.0 | 61.0% |
| Culiseta | 11.0 | 1.7% |
| Other | 6.0 | 0.9% |

FC-015

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 Stuart and Dorset

GPS: 40.5600827919436, -105.12403752654791


 Total number of trap/nights set:
 15.0

 Total number of mosquitoes collected:
 229.0

 Average mosquitoes per trap/night:
 15.3

 Average Culex per trap/night:
 3.0

| Aedes dorsalis | 8.0 | 3.5% |
|-------------------|-------|-------|
| Aedes hendersoni | 1.0 | 0.4% |
| Aedes increpitus | 13.0 | 5.7% |
| Aedes melanimon | 1.0 | 0.4% |
| Aedes trivittatus | 1.0 | 0.4% |
| Aedes vexans | 158.0 | 69.0% |
| Culex pipiens | 21.0 | 9.2% |
| Culex tarsalis | 24.0 | 10.5% |
| Culiseta inornata | 2.0 | 0.9% |
| | | |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 182.0 | 79.5% |
| Anopheles | 0.0 | 0.0% |
| Culex | 45.0 | 19.7% |
| Culiseta | 2.0 | 0.9% |
| Other | 0.0 | 0.0% |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 Edora Park

GPS: 40.565556701715764, -105.05251139402388

 Total number of trap/nights set:
 14.0

 Total number of mosquitoes collected:
 435.0

 Average mosquitoes per trap/night:
 31.1

 Average Culex per trap/night:
 20.7

Species collected and abundance:

| Aedes dorsalis | 6.0 | 1.4% |
|---------------------------|-------|-------|
| Aedes hendersoni | 1.0 | 0.2% |
| Aedes increpitus | 36.0 | 8.3% |
| Aedes melanimon | 2.0 | 0.5% |
| Aedes vexans | 77.0 | 17.7% |
| Anopheles freeborni | 3.0 | 0.7% |
| Coquillettidia perturbans | 11.0 | 2.5% |
| Culex pipiens | 115.0 | 26.4% |
| Culex tarsalis | 175.0 | 40.2% |
| Culiseta inornata | 9.0 | 2.1% |
| | | |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 122.0 | 28.0% |
| Anopheles | 3.0 | 0.7% |
| Culex | 290.0 | 66.7% |
| Culiseta | 9.0 | 2.1% |
| Other | 11.0 | 2.5% |

FC-023

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

Location: Boltz

GPS: 40.54436640664932, -105.06441771984099


 Total number of trap/nights set:
 15.0

 Total number of mosquitoes collected:
 298.0

 Average mosquitoes per trap/night:
 19.9

 Average Culex per trap/night:
 8.6

| Aedes dorsalis | 94.0 | 31.5% |
|-------------------|------|-------|
| Aedes increpitus | 1.0 | 0.3% |
| Aedes melanimon | 3.0 | 1.0% |
| Aedes vexans | 65.0 | 21.8% |
| Culex pipiens | 54.0 | 18.1% |
| Culex tarsalis | 75.0 | 25.2% |
| Culiseta inornata | 6.0 | 2.0% |
| | | |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 163.0 | 54.7% |
| Anopheles | 0.0 | 0.0% |
| Culex | 129.0 | 43.3% |
| Culiseta | 6.0 | 2.0% |
| Other | 0.0 | 0.0% |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 3001 San Luis

GPS: 40.546510068398455, -105.03359008580445

 Total number of trap/nights set:
 15.0


 Total number of mosquitoes collected:
 441.0

 Average mosquitoes per trap/night:
 29.4

 Average Culex per trap/night:
 14.9

Species collected and abundance:

| Aedes cinereus | 2.0 | 0.5% |
|---------------------------|-------|-------|
| Aedes dorsalis | 13.0 | 2.9% |
| Aedes increpitus | 44.0 | 10.0% |
| Aedes melanimon | 2.0 | 0.5% |
| Aedes trivittatus | 1.0 | 0.2% |
| Aedes vexans | 152.0 | 34.5% |
| Coquillettidia perturbans | 1.0 | 0.2% |
| Culex pipiens | 134.0 | 30.4% |
| Culex tarsalis | 89.0 | 20.2% |
| Culiseta inornata | 3.0 | 0.7% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 214.0 | 48.5% |
| Anopheles | 0.0 | 0.0% |
| Culex | 223.0 | 50.6% |
| Culiseta | 3.0 | 0.7% |
| Other | 1.0 | 0.2% |

FC-029

Season: 05/01/2024 - 09/17/2024
Trap Type: CDC Light Trap

Location: Ben's Park

GPS: 40.51136844640155, -105.07185079157352

Total number of trap/nights set: 15.0

Total number of mosquitoes collected: 378.0

Average mosquitoes per trap/night: 25.2

Average Culex per trap/night: 10.6

| | | 8.5% |
|-------------------|-------|-------|
| Aedes increpitus | 1.0 | 0.3% |
| Aedes melanimon | 2.0 | 0.5% |
| Aedes vexans | 182.0 | 48.1% |
| Culex pipiens | 52.0 | 13.8% |
| Culex tarsalis | 107.0 | 28.3% |
| Culiseta inornata | 2.0 | 0.5% |
| Trap Stolen | 0.0 | 0.0% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 217.0 | 57.4% |
| Anopheles | 0.0 | 0.0% |
| Culex | 159.0 | 42.1% |
| Culiseta | 2.0 | 0.5% |
| Other | 0.0 | 0.0% |

FC-029gr

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 Gravid Trap

 Location:
 Ben's Park Gravid

GPS: 40.51136309344342, -105.07115509361029

 Total number of trap/nights set:
 14.0

 Total number of mosquitoes collected:
 293.0


 Average mosquitoes per trap/night:
 20.9


 Average Culex per trap/night:
 20.9

Species collected and abundance:

 Aedes vexans
 1.0
 0.3%

 Culex pipiens
 292.0
 99.7%





FC-031

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 Willow Springs

GPS: 40.50608996726513, -105.03941986709833


 Total number of trap/nights set:
 15.0

 Total number of mosquitoes collected:
 1,805.0

 Average mosquitoes per trap/night:
 120.3

 Average Culex per trap/night:
 39.2

| Aedes dorsalis | 402.0 | 22.3% |
|--------------------|-------|-------|
| Aedes hendersoni | 1.0 | 0.1% |
| Aedes melanimon | 51.0 | 2.8% |
| Aedes nigromaculis | 2.0 | 0.1% |
| Aedes vexans | 746.0 | 41.3% |
| Culex pipiens | 57.0 | 3.2% |
| Culex tarsalis | 531.0 | 29.4% |
| Culiseta inornata | 15.0 | 0.8% |



| Genus Proportions: | | |
|--------------------|---------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 1,202.0 | 66.6% |
| Anopheles | 0.0 | 0.0% |
| Culex | 588.0 | 32.6% |
| Culiseta | 15.0 | 0.8% |
| Other | 0.0 | 0.0% |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 Country Club

GPS: 40.62669924225406, -105.0521868467331

| Total number of trap/nights set: | 15.0 |
|---------------------------------------|---------|
| Total number of mosquitoes collected: | 2,303.0 |
| Average mosquitoes per trap/night: | 153.5 |
| Average Culex per trap/night: | 10.7 |

Species collected and abundance:

| Aedes dorsalis | 169.0 | 7.3% |
|---------------------|---------|-------|
| Aedes increpitus | 9.0 | 0.4% |
| Aedes melanimon | 434.0 | 18.8% |
| Aedes trivittatus | 2.0 | 0.1% |
| Aedes vexans | 1,200.0 | 52.1% |
| Anopheles freeborni | 316.0 | 13.7% |
| Culex pipiens | 2.0 | 0.1% |
| Culex tarsalis | 159.0 | 6.9% |
| Culiseta inornata | 12.0 | 0.5% |



| Genus Proportions: | | |
|--------------------|---------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 1,814.0 | 78.8% |
| Anopheles | 316.0 | 13.7% |
| Culex | 161.0 | 7.0% |
| Culiseta | 12.0 | 0.5% |
| Other | 0.0 | 0.0% |

FC-036

 Season:
 05/01/2024 - 09/17/2024


 Trap Type:
 CDC Light Trap

ocation: Hemlock

GPS: 40.60076561609448, -105.07983673363923

| Total number of trap/nights set: | 15.0 |
|---------------------------------------|---------|
| Total number of mosquitoes collected: | 2,179.0 |
| Average mosquitoes per trap/night: | 145.3 |
| Average Culex per trap/night: | 37.2 |

| Aedes dorsalis | 125.0 | 5.7% |
|---------------------------|---------|-------|
| Aedes increpitus | 197.0 | 9.0% |
| Aedes melanimon | 33.0 | 1.5% |
| Aedes vexans | 1,195.0 | 54.8% |
| Anopheles freeborni | 9.0 | 0.4% |
| Coquillettidia perturbans | 25.0 | 1.1% |
| Culex pipiens | 240.0 | 11.0% |
| Culex tarsalis | 318.0 | 14.6% |
| Culiseta inornata | 37.0 | 1.7% |



| Genus Proportions: | | |
|--------------------|---------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 1,550.0 | 71.1% |
| Anopheles | 9.0 | 0.4% |
| Culex | 558.0 | 25.6% |
| Culiseta | 37.0 | 1.7% |
| Other | 25.0 | 1.1% |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

Location: Chelsea Ridge

GPS: 40.51667989969775, -105.09808011353017

 Total number of trap/nights set:
 15.0


 Total number of mosquitoes collected:
 970.0

 Average mosquitoes per trap/night:
 64.7

 Average Culex per trap/night:
 14.0

Species collected and abundance:

| Aedes dorsalis | 6.0 | 0.6% |
|---------------------|-------|-------|
| Aedes hendersoni | 1.0 | 0.1% |
| Aedes increpitus | 88.0 | 9.1% |
| Aedes melanimon | 6.0 | 0.6% |
| Aedes trivittatus | 1.0 | 0.1% |
| Aedes vexans | 648.0 | 66.8% |
| Anopheles freeborni | 1.0 | 0.1% |
| Culex pipiens | 41.0 | 4.2% |
| Culex tarsalis | 169.0 | 17.4% |
| Culiseta inornata | 9.0 | 0.9% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 750.0 | 77.3% |
| Anopheles | 1.0 | 0.1% |
| Culex | 210.0 | 21.6% |
| Culiseta | 9.0 | 0.9% |
| Other | 0.0 | 0.0% |

FC-038


Season: 05/01/2024 - 09/17/2024

Trap Type: CDC Light Trap
Location: Lochside Lane

GPS: 40.59907046206435, -105.00655643641949

| Total number of trap/nights set: | 15.0 |
|---------------------------------------|---------|
| Total number of mosquitoes collected: | 1,424.0 |
| Average mosquitoes per trap/night: | 94.9 |
| Average Culex per trap/night: | 32.1 |

| Aedes dorsalis | 56.0 | 3.9% |
|---------------------------|-------|-------|
| Aedes increpitus | 18.0 | 1.3% |
| Aedes melanimon | 128.0 | 9.0% |
| Aedes trivittatus | 3.0 | 0.2% |
| Aedes vexans | 709.0 | 49.8% |
| Anopheles freeborni | 4.0 | 0.3% |
| Coquillettidia perturbans | 1.0 | 0.1% |
| Culex pipiens | 53.0 | 3.7% |
| Culex tarsalis | 428.0 | 30.1% |
| Culiseta inornata | 24.0 | 1.7% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 914.0 | 64.2% |
| Anopheles | 4.0 | 0.3% |
| Culex | 481.0 | 33.8% |
| Culiseta | 24.0 | 1.7% |
| Other | 1.0 | 0.1% |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 Fossil Creek South

GPS: 40.4808101171896, -105.03934007138014

 Total number of trap/nights set:
 15.0


 Total number of mosquitoes collected:
 708.0


 Average mosquitoes per trap/night:
 47.2

 Average Culex per trap/night:
 8.7

Species collected and abundance:

| Aedes dorsalis | 323.0 | 45.6% |
|---------------------|-------|-------|
| Aedes increpitus | 1.0 | 0.1% |
| Aedes melanimon | 25.0 | 3.5% |
| Aedes vexans | 183.0 | 25.8% |
| Anopheles freeborni | 5.0 | 0.7% |
| Culex pipiens | 3.0 | 0.4% |
| Culex tarsalis | 127.0 | 17.9% |
| Culiseta inornata | 41.0 | 5.8% |





FC-040

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 Redwood

GPS: 40.60307751389784, -105.06707444787025


 Total number of trap/nights set:
 15.0

 Total number of mosquitoes collected:
 750.0

 Average mosquitoes per trap/night:
 50.0

 Average Culex per trap/night:
 17.8

| Aedes dorsalis | 96.0 | 12.8% |
|---------------------------|-------|-------|
| Aedes increpitus | 5.0 | 0.7% |
| Aedes melanimon | 27.0 | 3.6% |
| Aedes vexans | 325.0 | 43.3% |
| Anopheles freeborni | 1.0 | 0.1% |
| Coquillettidia perturbans | 24.0 | 3.2% |
| Culex pipiens | 87.0 | 11.6% |
| Culex tarsalis | 180.0 | 24.0% |
| Culiseta inornata | 5.0 | 0.7% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 453.0 | 60.4% |
| Anopheles | 1.0 | 0.1% |
| Culex | 267.0 | 35.6% |
| Culiseta | 5.0 | 0.7% |
| Other | 24.0 | 3.2% |

FC-040gr

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 Gravid Trap

 Location:
 Redwood Gravid

GPS: 40.60309558721682, -105.06650783121586

 Total number of trap/nights set:
 15.0


 Total number of mosquitoes collected:
 1,037.0

 Average mosquitoes per trap/night:
 69.1

 Average Culex per trap/night:
 68.3

Species collected and abundance:

| Aedes increpitus | 2.0 | 0.2% |
|-------------------|---------|-------|
| Aedes vexans | 9.0 | 0.9% |
| Culex pipiens | 1,016.0 | 98.0% |
| Culex tarsalis | 8.0 | 0.8% |
| Culiseta inornata | 2.0 | 0.2% |



| Genus Proportions: | | |
|--------------------|---------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 11.0 | 1.1% |
| Anopheles | 0.0 | 0.0% |
| Culex | 1,024.0 | 98.7% |
| Culiseta | 2.0 | 0.2% |
| Other | 0.0 | 0.0% |

FC-041

 Season:
 05/01/2024 - 09/17/2024


 Trap Type:
 CDC Light Trap

ocation: Fishback

GPS: 40.58794002990494, -105.10486006736755

| Total number of trap/nights set: | 14.0 |
|---------------------------------------|-------|
| Total number of mosquitoes collected: | 266.0 |
| Average mosquitoes per trap/night: | 19.0 |
| Average Culex per trap/night: | 10.0 |

| Aedes dorsalis | 1.0 | 0.4% |
|-------------------|-------|-------|
| Aedes hendersoni | 2.0 | 0.8% |
| Aedes melanimon | 6.0 | 2.3% |
| Aedes vexans | 115.0 | 43.2% |
| Culex pipiens | 88.0 | 33.1% |
| Culex tarsalis | 52.0 | 19.5% |
| Culiseta inornata | 2.0 | 0.8% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 124.0 | 46.6% |
| Anopheles | 0.0 | 0.0% |
| Culex | 140.0 | 52.6% |
| Culiseta | 2.0 | 0.8% |
| Other | 0.0 | 0.0% |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 Westshore Ct

GPS: 40.52954284354251, -105.0651342049241

 Total number of trap/nights set:
 15.0


 Total number of mosquitoes collected:
 340.0

 Average mosquitoes per trap/night:
 22.7

 Average Culex per trap/night:
 11.8

Species collected and abundance:

| Aedes cinereus | 1.0 | 0.3% |
|--------------------|-------|-------|
| Aedes dorsalis | 33.0 | 9.7% |
| Aedes increpitus | 3.0 | 0.9% |
| Aedes melanimon | 3.0 | 0.9% |
| Aedes nigromaculis | 1.0 | 0.3% |
| Aedes vexans | 120.0 | 35.3% |
| Culex pipiens | 75.0 | 22.1% |
| Culex tarsalis | 102.0 | 30.0% |
| Culiseta inornata | 2.0 | 0.6% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 161.0 | 47.4% |
| Anopheles | 0.0 | 0.0% |
| Culex | 177.0 | 52.1% |
| Culiseta | 2.0 | 0.6% |
| Other | 0.0 | 0.0% |

FC-047

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 Keeneland And Twin Oak

GPS: 40.51511286336567, -105.0528560578823


Total number of trap/nights set: 14.0

Total number of mosquitoes collected: 510.0

Average mosquitoes per trap/night: 36.4

Average Culex per trap/night: 20.7

| Aedes dorsalis | 12.0 | 2.49 |
|---------------------|-------|-------|
| Aedes increpitus | 3.0 | 0.69 |
| Aedes melanimon | 2.0 | 0.49 |
| Aedes vexans | 199.0 | 39.0% |
| Anopheles freeborni | 1.0 | 0.29 |
| Culex pipiens | 56.0 | 11.0% |
| Culex tarsalis | 234.0 | 45.99 |
| Culiseta inornata | 3.0 | 0.69 |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 216.0 | 42.4% |
| Anopheles | 1.0 | 0.2% |
| Culex | 290.0 | 56.9% |
| Culiseta | 3.0 | 0.6% |
| Other | 0.0 | 0.0% |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 Casa Grande and Downing

GPS: 40.573239949533054, -105.13949006795883

 Total number of trap/nights set:
 13.0


 Total number of mosquitoes collected:
 225.0

 Average mosquitoes per trap/night:
 17.3

 Average Culex per trap/night:
 9.5

Species collected and abundance:

| Aedes dorsalis | 4.0 | 1.8% |
|-------------------|------|-------|
| Aedes increpitus | 4.0 | 1.8% |
| Aedes melanimon | 2.0 | 0.9% |
| Aedes vexans | 88.0 | 39.1% |
| Culex pipiens | 81.0 | 36.0% |
| Culex tarsalis | 43.0 | 19.1% |
| Culiseta inornata | 3.0 | 1.3% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 98.0 | 43.6% |
| Anopheles | 0.0 | 0.0% |
| Culex | 124.0 | 55.1% |
| Culiseta | 3.0 | 1.3% |
| Other | 0.0 | 0.0% |

FC-050

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 Golden Meadows Ditch

 GPS:
 40.52954004037219, -105.05033008754252


 Total number of trap/nights set:
 15,0

 Total number of mosquitoes collected:
 161.0

 Average mosquitoes per trap/night:
 10.7

 Average Culex per trap/night:
 7.5

| Aedes dorsalis | 3.0 | 1.9% |
|-------------------|------|-------|
| Aedes melanimon | 4.0 | 2.5% |
| Aedes vexans | 40.0 | 24.8% |
| Culex pipiens | 64.0 | 39.8% |
| Culex tarsalis | 49.0 | 30.4% |
| Culiseta inornata | 1.0 | 0.6% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 47.0 | 29.2% |
| Anopheles | 0.0 | 0.0% |
| Culex | 113.0 | 70.2% |
| Culiseta | 1.0 | 0.6% |
| Other | 0.0 | 0.0% |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 603 Gilgalad Way

GPS: 40.5616299281039, -105.08703008294106

 Total number of trap/nights set:
 15.0


 Total number of mosquitoes collected:
 1,418.0

 Average mosquitoes per trap/night:
 94.5

 Average Culex per trap/night:
 4.4

Species collected and abundance:

| Aedes cinereus | 3.0 | 0.2% |
|---------------------|---------|-------|
| Aedes dorsalis | 1.0 | 0.1% |
| Aedes hendersoni | 1.0 | 0.1% |
| Aedes increpitus | 24.0 | 1.7% |
| Aedes melanimon | 6.0 | 0.4% |
| Aedes vexans | 1,281.0 | 90.3% |
| Anopheles freeborni | 1.0 | 0.1% |
| Culex pipiens | 23.0 | 1.6% |
| Culex tarsalis | 43.0 | 3.0% |
| Culiseta inornata | 35.0 | 2.5% |



| Genus Proportions: | | |
|--------------------|---------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 1,316.0 | 92.8% |
| Anopheles | 1.0 | 0.1% |
| Culex | 66.0 | 4.7% |
| Culiseta | 35.0 | 2.5% |
| Other | 0.0 | 0.0% |

FC-053

Season: 05/01/2024 - 09/17/2024
Trap Type: CDC Light Trap

Trap Type: CDC Light Trap

Location: Egret and Rookery

GPS: 40.4994200898768, -105.01182999461888


 Total number of trap/nights set:
 14.0

 Total number of mosquitoes collected:
 541.0

 Average mosquitoes per trap/night:
 38.6

 Average Culex per trap/night:
 24.0

| Aedes dorsalis | 28.0 | 5.2% |
|---------------------------|-------|-------|
| Aedes melanimon | 20.0 | 3.7% |
| Aedes vexans | 146.0 | 27.0% |
| Anopheles freeborni | 4.0 | 0.7% |
| Coquillettidia perturbans | 1.0 | 0.2% |
| Culex pipiens | 19.0 | 3.5% |
| Culex tarsalis | 317.0 | 58.6% |
| Culiseta inornata | 6.0 | 1.1% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 194.0 | 35.9% |
| Anopheles | 4.0 | 0.7% |
| Culex | 336.0 | 62.1% |
| Culiseta | 6.0 | 1.1% |
| Other | 1.0 | 0.2% |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 737 Parliament

GPS: 40.49997995372379, -105.06319999694824

 Total number of trap/nights set:
 15.0


 Total number of mosquitoes collected:
 285.0

 Average mosquitoes per trap/night:
 19.0

 Average Culex per trap/night:
 3.3

Species collected and abundance:

| Aedes dorsalis | 26.0 | 9.1% |
|-------------------|-------|-------|
| Aedes increpitus | 1.0 | 0.4% |
| Aedes melanimon | 2.0 | 0.7% |
| Aedes vexans | 201.0 | 70.5% |
| Culex pipiens | 2.0 | 0.7% |
| Culex tarsalis | 47.0 | 16.5% |
| Culiseta inornata | 6.0 | 2.1% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 230.0 | 80.7% |
| Anopheles | 0.0 | 0.0% |
| Culex | 49.0 | 17.2% |
| Culiseta | 6.0 | 2.1% |
| Other | 0.0 | 0.0% |

FC-057

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

Location: Registry Ridge

GPS: 40.48438997678501, -105.10523993521929


 Total number of trap/nights set:
 15.0

 Total number of mosquitoes collected:
 332.0

 Average mosquitoes per trap/night:
 22.1

 Average Culex per trap/night:
 2.2

| Aedes dorsalis | 180.0 | 54.2% |
|-------------------|-------|-------|
| Aedes increpitus | 1.0 | 0.3% |
| Aedes melanimon | 5.0 | 1.5% |
| Aedes vexans | 106.0 | 31.9% |
| Culex pipiens | 6.0 | 1.8% |
| Culex tarsalis | 27.0 | 8.1% |
| Culiseta inornata | 7.0 | 2.1% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 292.0 | 88.0% |
| Anopheles | 0.0 | 0.0% |
| Culex | 33.0 | 9.9% |
| Culiseta | 7.0 | 2.1% |
| Other | 0.0 | 0.0% |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 Spring Creek Trail-- Michener Dr

 GPS:
 40.54883989155028, -105.12533001601697

 Total number of trap/nights set:
 15.0


 Total number of mosquitoes collected:
 281.0

 Average mosquitoes per trap/night:
 18.7

 Average Culex per trap/night:
 3.4

Species collected and abundance:

| Aedes dorsalis | 3.0 | 1.1% |
|-------------------|-------|-------|
| Aedes increpitus | 68.0 | 24.2% |
| Aedes melanimon | 1.0 | 0.4% |
| Aedes vexans | 149.0 | 53.0% |
| Culex pipiens | 22.0 | 7.8% |
| Culex tarsalis | 29.0 | 10.3% |
| Culiseta inornata | 9.0 | 3.2% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 221.0 | 78.6% |
| Anopheles | 0.0 | 0.0% |
| Culex | 51.0 | 18.1% |
| Culiseta | 9.0 | 3.2% |
| Other | 0.0 | 0.0% |

FC-059

Season: 05/01/2024 - 09/17/2024
Trap Type: CDC Light Trap

 Location:
 Springwood and Lochwood

 GPS:
 40.5426000124595, -105.0460398942232


 Total number of trap/nights set:
 15.0

 Total number of mosquitoes collected:
 135.0

 Average mosquitoes per trap/night:
 9.0

 Average Culex per trap/night:
 3.5

| Aedes dorsalis | 7.0 | 5.2% |
|-------------------|------|-------|
| Aedes increpitus | 1.0 | 0.7% |
| Aedes vexans | 65.0 | 48.1% |
| Culex pipiens | 15.0 | 11.1% |
| Culex tarsalis | 38.0 | 28.1% |
| Culiseta inornata | 9.0 | 6.7% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 73.0 | 54.1% |
| Anopheles | 0.0 | 0.0% |
| Culex | 53.0 | 39.3% |
| Culiseta | 9.0 | 6.7% |
| Other | 0.0 | 0.0% |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 808 Ponderosa

GPS: 40.548700028018544, -105.12018989771605

 Total number of trap/nights set:
 15.0


 Total number of mosquitoes collected:
 152.0


 Average mosquitoes per trap/night:
 10.1

 Average Culex per trap/night:
 7.1

Species collected and abundance:

| Aedes dorsalis | 4.0 | 2.69 |
|---------------------|------|-------|
| Aedes increpitus | 1.0 | 0.79 |
| Aedes melanimon | 2.0 | 1.39 |
| Aedes vexans | 35.0 | 23.09 |
| Anopheles freeborni | 1.0 | 0.79 |
| Culex pipiens | 63.0 | 41.49 |
| Culex tarsalis | 44.0 | 28.99 |
| Culiseta inornata | 2.0 | 1.39 |
| | | |





FC-061

 Season:
 05/01/2024 - 09/17/2024


 Trap Type:
 CDC Light Trap

Location: Holley Plant Research Center

GPS: 40.56879993041706, -105.09164985269308

| Total number of trap/nights set: | 15.0 |
|---------------------------------------|---------|
| Total number of mosquitoes collected: | 1,366.0 |
| Average mosquitoes per trap/night: | 91.1 |
| Average Culex per trap/night: | 27.2 |

| Aedes dorsalis | 17.0 | 1.2% |
|---------------------------|-------|-------|
| Aedes hendersoni | 7.0 | 0.5% |
| Aedes increpitus | 168.0 | 12.3% |
| Aedes melanimon | 7.0 | 0.5% |
| Aedes vexans | 738.0 | 54.0% |
| Coquillettidia perturbans | 2.0 | 0.1% |
| Culex pipiens | 196.0 | 14.3% |
| Culex tarsalis | 212.0 | 15.5% |
| Culiseta incidens | 4.0 | 0.3% |
| Culiseta inornata | 15.0 | 1.1% |
| | | |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 937.0 | 68.6% |
| Anopheles | 0.0 | 0.0% |
| Culex | 408.0 | 29.9% |
| Culiseta | 19.0 | 1.4% |
| Other | 2.0 | 0.1% |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap


 Location:
 Water's Edge at Blue Mesa

GPS: 40.542779889722866, -105.08740995079279

| Total number of trap/nights set: | 14.0 |
|---------------------------------------|-------|
| Total number of mosquitoes collected: | 227.0 |
| Average mosquitoes per trap/night: | 16.2 |
| Average Culex per trap/night: | 5.9 |

Species collected and abundance:

| Aedes dorsalis | 24.0 | 10.6% |
|-------------------|-------|-------|
| Aedes hendersoni | 1.0 | 0.4% |
| Aedes idahoensis | 1.0 | 0.4% |
| Aedes increpitus | 10.0 | 4.4% |
| Aedes trivittatus | 2.0 | 0.9% |
| Aedes vexans | 105.0 | 46.3% |
| Culex pipiens | 45.0 | 19.8% |
| Culex tarsalis | 37.0 | 16.3% |
| Culiseta inornata | 2.0 | 0.9% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 143.0 | 63.0% |
| Anopheles | 0.0 | 0.0% |
| Culex | 82.0 | 36.1% |
| Culiseta | 2.0 | 0.9% |
| Other | 0.0 | 0.0% |

FC-063

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

Location: Red Fox Meadows

GPS: 40.565679974786285, -105.10485000908375

| Total number of trap/nights set: | 15.0 |
|---------------------------------------|---------|
| Total number of mosquitoes collected: | 4,759.0 |
| Average mosquitoes per trap/night: | 317.3 |
| Average Culex per trap/night: | 2.8 |

| Aedes dorsalis | 23.0 | 0.5% |
|---------------------|---------|-------|
| Aedes increpitus | 3,532.0 | 74.2% |
| Aedes melanimon | 2.0 | 0.0% |
| Aedes trivittatus | 3.0 | 0.1% |
| Aedes vexans | 1,148.0 | 24.1% |
| Anopheles freeborni | 2.0 | 0.0% |
| Culex pipiens | 19.0 | 0.4% |
| Culex tarsalis | 23.0 | 0.5% |
| Culiseta inornata | 7.0 | 0.1% |
| | | |



| Genus Proportions: | | |
|--------------------|---------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 4,708.0 | 98.9% |
| Anopheles | 2.0 | 0.0% |
| Culex | 42.0 | 0.9% |
| Culiseta | 7.0 | 0.1% |
| Other | 0.0 | 0.0% |

FC-063gr

Season: 05/01/2024 - 09/17/2024
Trap Type: Gravid Trap

Location: Red Fox Meadows Gravid

GPS: 40.565679974786285, -105.10504983365536

 Total number of trap/nights set:
 15.0


 Total number of mosquitoes collected:
 640.0

 Average mosquitoes per trap/night:
 42.7

 Average Culex per trap/night:
 25.6

Species collected and abundance:

| Aedes increpitus | 163.0 | 25.5% |
|------------------|-------|-------|
| Aedes melanimon | 13.0 | 2.0% |
| Aedes vexans | 80.0 | 12.5% |
| Culex pipiens | 382.0 | 59.7% |
| Culex tarsalis | 2.0 | 0.3% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 256.0 | 40.0% |
| Anopheles | 0.0 | 0.0% |
| Culex | 384.0 | 60.0% |
| Culiseta | 0.0 | 0.0% |
| Other | 0.0 | 0.0% |

FC-064

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

Location: West Chase

GPS: 40.49867997125427, -105.0298001244664

 Total number of trap/nights set:
 15.0

 Total number of mosquitoes collected:
 1,145.0

 Average mosquitoes per trap/night:
 76.3

 Average Culex per trap/night:
 22.6

| Aedes cinereus | 102.0 | 8.9% |
|---------------------|-------|-------|
| Aedes dorsalis | 25.0 | 2.2% |
| Aedes hendersoni | 1.0 | 0.1% |
| Aedes increpitus | 13.0 | 1.1% |
| Aedes melanimon | 37.0 | 3.2% |
| Aedes trivittatus | 1.0 | 0.1% |
| Aedes vexans | 622.0 | 54.3% |
| Anopheles freeborni | 1.0 | 0.1% |
| Culex pipiens | 79.0 | 6.9% |
| Culex tarsalis | 260.0 | 22.7% |
| Culiseta inornata | 4.0 | 0.3% |
| | | |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 801.0 | 70.0% |
| Anopheles | 1.0 | 0.1% |
| Culex | 339.0 | 29.6% |
| Culiseta | 4.0 | 0.3% |
| Other | 0.0 | 0.0% |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 Prospect Ponds

GPS: 40.55846837302044, -105.02281665802002

 Total number of trap/nights set:
 15.0


 Total number of mosquitoes collected:
 3,831.0

 Average mosquitoes per trap/night:
 255.4

 Average Culex per trap/night:
 56.6

Species collected and abundance:

| Aedes dorsalis | 162.0 | 4.2% |
|---------------------------|---------|-------|
| Aedes hendersoni | 8.0 | 0.2% |
| Aedes increpitus | 233.0 | 6.1% |
| Aedes melanimon | 66.0 | 1.7% |
| Aedes nigromaculis | 2.0 | 0.1% |
| Aedes vexans | 1,400.0 | 36.5% |
| Anopheles freeborni | 126.0 | 3.3% |
| Coquillettidia perturbans | 928.0 | 24.2% |
| Culex pipiens | 343.0 | 9.0% |
| Culex tarsalis | 506.0 | 13.2% |
| Culiseta inornata | 57.0 | 1.5% |



| Genus Proportions: | | |
|--------------------|---------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 1,871.0 | 48.8% |
| Anopheles | 126.0 | 3.3% |
| Culex | 849.0 | 22.2% |
| Culiseta | 57.0 | 1.5% |
| Other | 928.0 | 24.2% |

FC-066gr

Season: 05/01/2024 - 09/17/2024

Trap Type: Gravid Trap

Location: Prospect Ponds Gravid

GPS: 40.55830305717813, -105.02268858253954


 Total number of trap/nights set:
 15.0

 Total number of mosquitoes collected:
 291.0

 Average mosquitoes per trap/night:
 19.4

 Average Culex per trap/night:
 13.3

| Aedes dorsalis | 3.0 | 1.0% |
|---------------------------|-------|-------|
| Aedes increpitus | 4.0 | 1.4% |
| Aedes vexans | 42.0 | 14.4% |
| Anopheles freeborni | 35.0 | 12.0% |
| Coquillettidia perturbans | 2.0 | 0.7% |
| Culex pipiens | 195.0 | 67.0% |
| Culex tarsalis | 4.0 | 1.4% |
| Culiseta inornata | 6.0 | 2.1% |



| Genus Proportions: | | | |
|--------------------|--------|------------------|--|
| Genus | Number | Percent of Total | |
| Aedes/Ochlerotatus | 49.0 | 16.8% | |
| Anopheles | 35.0 | 12.0% | |
| Culex | 199.0 | 68.4% | |
| Culiseta | 6.0 | 2.1% | |
| Other | 2.0 | 0.7% | |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 Poudre River Trail

GPS: 40.57749479750878, -105.05568780004978

 Total number of trap/nights set:
 15.0

 Total number of mosquitoes collected:
 802.0

 Average mosquitoes per trap/night:
 53.5

 Average Culex per trap/night:
 4.9

Species collected and abundance:

| Aedes dorsalis | 41.0 | 5.1% |
|---------------------------|-------|-------|
| Aedes increpitus | 105.0 | 13.1% |
| Aedes melanimon | 24.0 | 3.0% |
| Aedes nigromaculis | 2.0 | 0.2% |
| Aedes vexans | 540.0 | 67.3% |
| Anopheles freeborni | 3.0 | 0.4% |
| Coquillettidia perturbans | 4.0 | 0.5% |
| Culex pipiens | 12.0 | 1.5% |
| Culex tarsalis | 62.0 | 7.7% |
| Culiseta inornata | 9.0 | 1.1% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 712.0 | 88.8% |
| Anopheles | 3.0 | 0.4% |
| Culex | 74.0 | 9.2% |
| Culiseta | 9.0 | 1.1% |
| Other | 4.0 | 0.5% |

FC-068

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 5029 Crest Dr

GPS: 40.516000124446485, -105.08547004312277


 Total number of trap/nights set:
 15.0

 Total number of mosquitoes collected:
 150.0

 Average mosquitoes per trap/night:
 10.0

 Average Culex per trap/night:
 2.5

| Aedes dorsalis | 3.0 | 2.0% |
|---------------------|-------|-------|
| Aedes melanimon | 2.0 | 1.3% |
| Aedes vexans | 100.0 | 66.7% |
| Anopheles freeborni | 5.0 | 3.3% |
| Culex pipiens | 8.0 | 5.3% |
| Culex tarsalis | 29.0 | 19.3% |
| Culiseta inornata | 3.0 | 2.0% |



| Genus Proportions: | | | |
|--------------------|--------|------------------|--|
| Genus | Number | Percent of Total | |
| Aedes/Ochlerotatus | 105.0 | 70.0% | |
| Anopheles | 5.0 | 3.3% | |
| Culex | 37.0 | 24.7% | |
| Culiseta | 3.0 | 2.0% | |
| Other | 0.0 | 0.0% | |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 Linden Lake Rd

GPS: 40.61451923788414, -105.05314875394106

 Total number of trap/nights set:
 14.0


 Total number of mosquitoes collected:
 1,158.0


 Average mosquitoes per trap/night:
 82.7

 Average Culex per trap/night:
 6.4

Species collected and abundance:

| Aedes dorsalis | 66.0 | 5.7% |
|---------------------------|-------|-------|
| Aedes increpitus | 46.0 | 4.0% |
| Aedes melanimon | 99.0 | 8.5% |
| Aedes trivittatus | 1.0 | 0.1% |
| Aedes vexans | 792.0 | 68.4% |
| Anopheles freeborni | 8.0 | 0.7% |
| Coquillettidia perturbans | 52.0 | 4.5% |
| Culex pipiens | 8.0 | 0.7% |
| Culex tarsalis | 82.0 | 7.1% |
| Culiseta inornata | 4.0 | 0.3% |





FC-071

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 Silvergate Rd

GPS: 40.52733008634481, -105.10821014642715


 Total number of trap/nights set:
 14.0

 Total number of mosquitoes collected:
 102.0

 Average mosquitoes per trap/night:
 7.3

 Average Culex per trap/night:
 1.4

| Aedes dorsalis | 7.0 | 6.9% |
|------------------|------|-------|
| Aedes increpitus | 5.0 | 4.9% |
| Aedes vexans | 70.0 | 68.6% |
| Culex pipiens | 10.0 | 9.8% |
| Culex tarsalis | 10.0 | 9.8% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 82.0 | 80.4% |
| Anopheles | 0.0 | 0.0% |
| Culex | 20.0 | 19.6% |
| Culiseta | 0.0 | 0.0% |
| Other | 0.0 | 0.0% |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap


 Location:
 422 Lake Dr

GPS: 40.569530106375986, -105.07116984575985

| Total number of trap/nights set: | 15.0 |
|---------------------------------------|-------|
| Total number of mosquitoes collected: | 442.0 |
| Average mosquitoes per trap/night: | 29.5 |
| Average Culex per trap/night: | 21.8 |

Species collected and abundance:

| Aedes dorsalis | 5.0 | 1.1% |
|-------------------|-------|-------|
| Aedes hendersoni | 2.0 | 0.5% |
| Aedes increpitus | 5.0 | 1.1% |
| Aedes melanimon | 5.0 | 1.1% |
| Aedes vexans | 95.0 | 21.5% |
| Culex pipiens | 139.0 | 31.4% |
| Culex tarsalis | 188.0 | 42.5% |
| Culiseta inornata | 3.0 | 0.7% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 112.0 | 25.3% |
| Anopheles | 0.0 | 0.0% |
| Culex | 327.0 | 74.0% |
| Culiseta | 3.0 | 0.7% |
| Other | 0.0 | 0.0% |

FC-073

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 118 S Grant

GPS: 40.58634003233039, -105.08899983018637


 Total number of trap/nights set:
 15.0

 Total number of mosquitoes collected:
 221.0

 Average mosquitoes per trap/night:
 14.7

 Average Culex per trap/night:
 9.1

| Aedes dorsalis | 2.0 | 0.9% |
|------------------|------|-------|
| Aedes hendersoni | 2.0 | 0.9% |
| Aedes melanimon | 1.0 | 0.5% |
| Aedes vexans | 80.0 | 36.2% |
| Culex pipiens | 87.0 | 39.4% |
| Culex tarsalis | 49.0 | 22.2% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 85.0 | 38.5% |
| Anopheles | 0.0 | 0.0% |
| Culex | 136.0 | 61.5% |
| Culiseta | 0.0 | 0.0% |
| Other | 0.0 | 0.0% |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 Rock Creek

GPS: 40.513749705683445, -105.00338338315488

 Total number of trap/nights set:
 14.0

 Total number of mosquitoes collected:
 759.0

 Average mosquitoes per trap/night:
 54.2

 Average Culex per trap/night:
 25.3

Species collected and abundance:

| Aedes cinereus | 1.0 | 0.1% |
|--------------------|-------|-------|
| Aedes dorsalis | 72.0 | 9.5% |
| Aedes increpitus | 2.0 | 0.3% |
| Aedes melanimon | 11.0 | 1.4% |
| Aedes nigromaculis | 2.0 | 0.3% |
| Aedes vexans | 310.0 | 40.8% |
| Culex pipiens | 25.0 | 3.3% |
| Culex tarsalis | 329.0 | 43.3% |
| Culiseta inornata | 7.0 | 0.9% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 398.0 | 52.4% |
| Anopheles | 0.0 | 0.0% |
| Culex | 354.0 | 46.6% |
| Culiseta | 7.0 | 0.9% |
| Other | 0.0 | 0.0% |

FC-075

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

Location: Sage Creek North

GPS: 40.511379917024655, -105.01986991614105


 Total number of trap/nights set:
 15.0

 Total number of mosquitoes collected:
 876.0

 Average mosquitoes per trap/night:
 58.4

 Average Culex per trap/night:
 23.7

| Aedes dorsalis | 10.0 | 1.1% |
|-------------------|-------|-------|
| Aedes increpitus | 3.0 | 0.3% |
| Aedes melanimon | 5.0 | 0.6% |
| Aedes vexans | 499.0 | 57.0% |
| Culex pipiens | 166.0 | 18.9% |
| Culex tarsalis | 189.0 | 21.6% |
| Culiseta inornata | 4.0 | 0.5% |
| | | |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 517.0 | 59.0% |
| Anopheles | 0.0 | 0.0% |
| Culex | 355.0 | 40.5% |
| Culiseta | 4.0 | 0.5% |
| Other | 0.0 | 0.0% |

FC-075gr

Season: 05/01/2024 - 09/17/2024
Trap Type: Gravid Trap

Location: Sage Creek North Gravid

GPS: 40.511379917024655, -105.02936996519566

 Total number of trap/nights set:
 14.0


 Total number of mosquitoes collected:
 175.0

 Average mosquitoes per trap/night:
 12.5

 Average Culex per trap/night:
 10.6

Species collected and abundance:

| Aedes increpitus | 1.0 | 0.6% |
|-------------------|-------|-------|
| Aedes vexans | 20.0 | 11.4% |
| Culex pipiens | 144.0 | 82.3% |
| Culex tarsalis | 5.0 | 2.9% |
| Culiseta inornata | 5.0 | 2.9% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 21.0 | 12.0% |
| Anopheles | 0.0 | 0.0% |
| Culex | 149.0 | 85.1% |
| Culiseta | 5.0 | 2.9% |
| Other | 0.0 | 0.0% |

FC-088gr

Season: 05/01/2024 - 09/17/2024

Trap Type: Gravid Trap

 Location:
 English Ranch Park Gravid

 GPS:
 40.53330994364783, -105.0305300205946


 Total number of trap/nights set:
 15.0

 Total number of mosquitoes collected:
 1,179.0

 Average mosquitoes per trap/night:
 78.6

 Average Culex per trap/night:
 77.3

| Aedes increpitus | 1.0 | 0.1% |
|---------------------|---------|-------|
| Aedes vexans | 15.0 | 1.3% |
| Anopheles freeborni | 1.0 | 0.1% |
| Culex pipiens | 1,152.0 | 97.7% |
| Culex tarsalis | 8.0 | 0.7% |
| Culiseta inornata | 2.0 | 0.2% |



| Genus Proportions: | | |
|--------------------|---------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 16.0 | 1.4% |
| Anopheles | 1.0 | 0.1% |
| Culex | 1,160.0 | 98.4% |
| Culiseta | 2.0 | 0.2% |
| Other | 0.0 | 0.0% |

FC-089gr

Season: 05/01/2024 - 09/17/2024

Trap Type: Gravid Trap

Average Culex per trap/night:

 Location:
 Kunz Ct. and Brook Dr. Gravid

 GPS:
 40.53631011931944, -105.10056015104055

 Total number of trap/nights set:
 15.0

 Total number of mosquitoes collected:
 279.0

 Average mosquitoes per trap/night:
 18.6

Species collected and abundance:

Culex pipiens 279.0 100.0%



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 0.0 | 0.0% |
| Anopheles | 0.0 | 0.0% |
| Culex | 279.0 | 100.0% |
| Culiseta | 0.0 | 0.0% |
| Other | 0.0 | 0.0% |

18.6

FC-090gr

Season: 05/01/2024 - 09/17/2024

Trap Type: Gravid Trap

 Location:
 Mountain Grandview Cemetery Gravid

 GPS:
 40.58195995773713, -105.111360065639

Total number of trap/nights set: 15.0

Total number of mosquitoes collected: 689.0

Average mosquitoes per trap/night: 45.9

Average Culex per trap/night: 45.7

| Aedes vexans | 2.0 | 0.3% |
|---------------------|-------|-------|
| Anopheles freeborni | 1.0 | 0.1% |
| Culex pipiens | 682.0 | 99.0% |
| Culex tarsalis | 3.0 | 0.4% |
| Culiseta inornata | 1.0 | 0.1% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 2.0 | 0.3% |
| Anopheles | 1.0 | 0.1% |
| Culex | 685.0 | 99.4% |
| Culiseta | 1.0 | 0.1% |
| Other | 0.0 | 0.0% |

FC-091gr

Season: 05/01/2024 - 09/17/2024
Trap Type: Gravid Trap

Location: PVH Gravid

GPS: 40.570520046200166, -105.05456998944281

Total number of trap/nights set: 15.0


Total number of mosquitoes collected: 126.0

Average mosquitoes per trap/night: 8.4

Average Culex per trap/night: 8.1

Species collected and abundance:

| Aedes dorsalis | 1.0 | 0.8% |
|-------------------|-------|-------|
| Aedes increpitus | 1.0 | 0.8% |
| Aedes vexans | 2.0 | 1.6% |
| Culex pipiens | 119.0 | 94.4% |
| Culex tarsalis | 2.0 | 1.6% |
| Culiseta inornata | 1.0 | 0.8% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 4.0 | 3.2% |
| Anopheles | 0.0 | 0.0% |
| Culex | 121.0 | 96.0% |
| Culiseta | 1.0 | 0.8% |
| Other | 0.0 | 0.0% |

FC-092gr

Season: 05/01/2024 - 09/17/2024

Trap Type: Gravid Trap

Location: Udall Natural Gravid

GPS: 40.587873321668276, -105.06916154175995


 Total number of trap/nights set:
 15.0

 Total number of mosquitoes collected:
 354.0

 Average mosquitoes per trap/night:
 23.6

 Average Culex per trap/night:
 23.3

| Aedes dorsalis | 1.0 | 0.3% |
|---------------------|-------|-------|
| Anopheles freeborni | 2.0 | 0.6% |
| Culex pipiens | 348.0 | 98.3% |
| Culex tarsalis | 2.0 | 0.6% |
| Culiseta inornata | 1.0 | 0.3% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 1.0 | 0.3% |
| Anopheles | 2.0 | 0.6% |
| Culex | 350.0 | 98.9% |
| Culiseta | 1.0 | 0.3% |
| Other | 0.0 | 0.0% |

 Season:
 05/01/2024 - 09/17/2024

 Trap Type:
 CDC Light Trap

 Location:
 Lopez Elementary

GPS: 40.53186000594886, -105.0882501527667


 Total number of trap/nights set:
 15.0

 Total number of mosquitoes collected:
 722.0

 Average mosquitoes per trap/night:
 48.1

 Average Culex per trap/night:
 6.5

| Aedes cinereus | 1.0 | 0.1% |
|-------------------|-------|-------|
| Aedes dorsalis | 18.0 | 2.5% |
| Aedes increpitus | 31.0 | 4.3% |
| Aedes vexans | 571.0 | 79.1% |
| Culex pipiens | 42.0 | 5.8% |
| Culex tarsalis | 56.0 | 7.8% |
| Culiseta inornata | 3.0 | 0.4% |



| Genus Proportions: | | |
|--------------------|--------|------------------|
| Genus | Number | Percent of Total |
| Aedes/Ochlerotatus | 621.0 | 86.0% |
| Anopheles | 0.0 | 0.0% |
| Culex | 98.0 | 13.6% |
| Culiseta | 3.0 | 0.4% |
| Other | 0.0 | 0.0% |

Appendix 2: Larimer County Adult Sample Pool Positive Results and Locations

2024 WNV Positive Pools

Start Date: 05/01/2024 End Date: 09/12/2024

| - | | | ۰ |
|---|---------|-------|---|
| | OTTO | 222.0 | ı |
| | 43 V E- | | ı |

| Trap Location | Total Positive Pools |
|---------------------|----------------------|
| 9th and DesMoines | 2 |
| Big Thompson | 0 |
| Boyd Lake | 0 |
| Cattail Pond | 3 |
| Cr 20 and 9 | 0 |
| Horseshoe Peninsula | 1 |

Total Positive: 6

2024 WNV Positive Pools

| - | | | | |
|---|----|----|---|-----|
| к | OF | th | | 11/ |
| | | ш | w | |

| Trap Location | Total Positive Pools |
|----------------|----------------------|
| Berthoud East | 0 |
| Berthoud North | 0 |
| Berthoud Park | 2 |
| Berthoud Point | 0 |
| Berthoud West | 0 |

Total Positive; 2

*** Please note that Larimer County Public Health in conjunction with the CDC conducted additional trapping this year in Berthoud and found an additional 10 positive pools not collected by VDCI.

2024 WNV Positive Pools

Start Date: 05/01/2024 **End Date:** 09/12/2024

Fort Collins

| Trap | Location | Total Positive Pools | |
|---------------------|-------------------|----------------------|--|
| 118 S Grant | | 0 | |
| 3001 San Luis | | 0 | |
| 422 Lake Dr | | 2 | |
| 5029 Crest Dr | | 0 | |
| 603 Gilgalad Way | | 1 | |
| 737 Parliament | | 0 | |
| 808 Ponderosa | | 1 | |
| Ben's Park | | 0 | |
| Ben's Park Gravid | | 1 | |
| Big Horn | | 1 | |
| Boltz | | 0 | |
| Casa Grande and Do | owning | 0 | |
| Chelsea Ridge | | 0 | |
| Country Club | | 0 | |
| Edora Park | | 2 | |
| Egret and Rookery | | 1 | |
| English Ranch Park | Gravid | 4 | |
| FC Visitor Center | | 0 | |
| Fishback | | 0 | |
| Fossil Creek South | | 1 | |
| Golden Current | | 0 | |
| Golden Meadows D | itch | 0 | |
| Hemlock | | 1 | |
| Holley Plant Resear | ch Center | 0 | |
| Keeneland And Twi | n Oak | 2 | |
| Kunz Ct, and Brook | Dr. Gravid | 1 | |
| Linden Lake Rd | | 1 | |
| Lochside Lane | | 2 | |
| Lopez Elementary | | 0 | |
| Magic Carpet | | 0 | |
| Mountain Grandvie | w Cemetery Gravid | 1 | |
| N. Linden | | 1 | |
| Poudre River Trail | | 0 | |

| Trap Location | Total Positive Pools | |
|--------------------------------|----------------------|--|
| Prospect Ponds | 2 | |
| Prospect Ponds Gravid | 3 | |
| PVH Gravid | 1 | |
| Red Fox Meadows | 0 | |
| Red Fox Meadows Gravid | 3 | |
| Redwood | 1 | |
| Redwood Gravid | 4 | |
| Registry Ridge | 0 | |
| Rock Creek | 1 | |
| Sage Creek North | 0 | |
| Sage Creek North Gravid | 1 | |
| Silvergate Rd | 0 | |
| Spring Creek Trail Michener Dr | 0 | |
| Springwood and Lochwood | 0 | |
| Stuart and Dorset | 0 | |
| Udall Natural Gravid | 3 | |
| Water's Edge at Blue Mesa | 0 | |
| West Chase | 2 | |
| Westshore Ct | 0 | |
| Willow Springs | 0 | |

Total Positive: 44